

UNDERSEA FEATURE NAME PROPOSAL

(See NOTE overleaf)

Ocean or Sea Arctic Ocean Name proposed NP-28 Channel

Coordinates : A - of midpoint or summit : Lat. **North 88°00'** , Long **45° West**
_____ kilometres in _____ direction from _____

and/or B - extremities (if linear feature) :

Lat. **85° 30' N** } to { Lat. **89° 30' N**
Long. **50° W** } } Long. **20° W**

Description (kind of feature) : submarine channel

Identifying or categorizing characteristics (shape, dimensions, total relief, least depth, steepness, etc.):

Three crossings exist and show width up to 20 km and depth up to 200 meter. All crossings show a higher channel levee on the east side, see figure 1 and 2 of attached publication in *Marine Geology*.

Associated features : _____

Chart reference :

Shown with name on chart No. **Not at present**

Shown but not named on chart No. _____

Not shown but within area covered by chart No. **IBCAO MAP OF THE ARCTIC OCEAN**

Reason for choice of name (if a person, state how associated with the feature to be named) :

Russian ice drift station North Pole 28 drifted across the feature in the summer of 1988 and recorded the seismic data that most clearly expresses the existence of the channel. Naming of the feature is a tribute to the effort of Russian ice stations to the exploration of the Arctic Ocean.

Discovery facts :

Date : **Summer 1988** by (individuals or ship) **Russian ice drift station "North Pole 28"**

By means of (equipment) : **Seismic reflection measurements**

Navigation used : **Celestial**

Estimated positional accuracy in nautical miles : _____

Description of survey (track spacing, line crossing, grid network, etc.) : **sea ice drift track**

Nature and repository of other survey activities (dredge samples, cores, magnetics, gravity, photographs, etc.) : **Unknown**

Supporting material : enclose, if possible, a sketch map of the survey area, profiles of the features, etc.,

with reference to prior publication, if any : **Scientific publication in *Marine Geology*, vol. 204, p. 317-324, 2004.**

Submitted by : **Prof. Yngve Kristoffersen**

Date : **April 06, 2004**

Address : **Department of Earth Science, University of Bergen, Allégaten 41, N-5007 Bergen, NORWAY**

Concurred in by (if applicable) : _____

Address : _____

National Authority (if any) : _____

Address : _____

NOTE : This form should be forwarded, when completed :

- a) **If the undersea feature is located in territorial waters :-**
to your "National Authority for Approval of Undersea Feature Names" or, if this does not exist or is not known, either to the International Hydrographic Bureau or to the Intergovernmental Oceanographic Commission (see addresses below);
- b) **If the undersea feature is located in international waters :-**
to the International Hydrographic Bureau or to the Intergovernmental Oceanographic Commission, at the following addresses :

International Hydrographic Bureau
4, quai Antoine 1^{er}
B.P. 445
MC 98011 MONACO CEDEX
Principality of MONACO
Fax: +377 93 10 81 40
E-mail: pac@ihb.mc


Intergovernmental Oceanographic Commission
UNESCO
Place de Fontenoy
75700 PARIS
FRANCE
Fax: +33 1 45 68 58 12
E-mail : info@unesco.org



ELSEVIER

Available online at www.sciencedirect.com

Marine Geology 3445 (2004) 1–8

www.elsevier.com/locate/margeo

A submarine fan in the Amundsen Basin, Arctic Ocean

Yngve Kristoffersen^{a,*}, Mikhail Y. Sorokin^{b,1}, Wilfried Jokat^c,
Oddny Svendsen^d

^a Department of Earth Science, University of Bergen, Allègaten 41, N-5007 Bergen, Norway

^b Polar Marine Geosurvey Expedition, St. Petersburg, Russia

^c Alfred Wegener Institute of Marine and Polar Research, Bremerhaven, Germany

^d Norwegian Petroleum Directorate, Postboks 300, N-4001 Stavanger, Norway

Received 5 June 2003; received in revised form 12 November 2003; accepted 9 December 2003

Abstract

Seismic reflection data from drifting ice stations and an icebreaker survey document the presence of a submarine fan along the east flank of the Lomonosov Ridge in the Amundsen Basin, Arctic Ocean. The fan extends from a source area at the North Greenland and Canadian Arctic continental margin to the North Pole. The fan probably developed in response to the combined effect of increased glacial sediment input during the Plio–Pleistocene and topographically confined transport of glacial debris in the deep sea passage between Lincoln Sea margin and the southern end of the Lomonosov Ridge.

© 2004 Published by Elsevier B.V.

Keywords: Arctic Ocean; Amundsen Basin; bottom features; submarine fan; glaci-marine sedimentation

1. Introduction

The deep basin between the North Pole and the Gakkel Ridge was outlined from spot soundings made by Russian scientists and appeared on a bathymetric chart of the Arctic Ocean published in 1954 (Burkhanov, 1957; Weber, 1983). Definition of distinct geomorphologic provinces became possible when continuous echo soundings first were obtained over the interior of the polar basin by the nuclear submarines *Nautilus* and *Skate* in


1958 (Dietz and Shumway, 1961). They observed an abyssal plain between Lomonosov Ridge and Gakkel Ridge, but also noted a modern abyssal seafloor slope of 0.5 m/km from the North Pole towards the Gakkel Ridge and expressed their curiosity about the origin of such a slope in an abyssal setting. The first continuous multi-channel seismic reflection profile across the Amundsen Basin obtained by *Polarstern* in 1991 along a similar track enabled Jokat et al. (1995) to present a seismic stratigraphic cross section of the Amundsen Basin. These observations combined with the seismic reflection results from the Russian ice station ‘North Pole 28’ (NP-28) and the US ice station ‘Arctic Research Laboratory Ice Station–II’ (Ar-lis-II) in 1965 (Ostenso and Wold, 1977) reveal the existence of a submarine fan extending from

¹ Deceased.

* Corresponding author. Tel.: +47-55583407;

Fax: +47-55583660.

E-mail address: yngve.kristoffersen@geo.uib.no (Y. Kristoffersen).



1 Fig. 1. Map of the Amundsen Basin with the locations of seismic profiles and observed crossings of submarine channels. Major
 2 sediment pathways indicated by a bold dark red line and the outline of the North Pole Submarine Fan by dark brown colour.
 3 Tributary paths from the Canadian–Greenland continental slope inferred from the position of bathymetric downslope depres-
 4 sions. Bathymetry from International Bathymetric Map of the Arctic Ocean (Jakobsson et al., 2001).


48 a source area on the Lincoln Sea margin (Greenland–
 49 Canadian shelf) to the North Pole (Fig. 1).
 50 We propose to name this fan the North Pole Sub-
 51 marine Fan, and the main channel, the NP-28
 52 Channel in recognition of the contribution of
 53 the Russian ice stations to exploration of the
 54 deep polar basin.

55 2. A submarine fan in the Amundsen Basin

56 We identify channel-levee complexes in all the
 57 three seismic reflection lines that presently exist of
 58 this area (Fig. 2): in the published line drawing of
 59 the data obtained by a sparker source and a single
 60 hydrophone during the drift of ice station Arlis-II
 61 (Ostenso and Wold, 1977), in the processed re-

62 cords from ice station NP-28 obtained with deto-
 63 nating caps and a geophone array on the ice, and
 64 in the multi-channel seismic records from *Polar-*
 65 *stern* using a 24-l airgun array and a 300-m-long,
 66 12-channel streamer (Fuetterer, 1992).


67 In the NP-28 crossing (Figs. 1 and 2), a single
 68 large channel, ca. 20 km wide and 120 m deep, is
 69 present at the foot of the Lomonosov Ridge. The
 70 elevated levee construction merges with the abys-
 71 sal plain more than 100 km away from the chan-
 72 nel. At least four relict channel positions are seen
 73 within the upper 500 milliseconds of levee stratig-
 74 raphy and indicate that the channel has migrated
 75 westward towards the foot of Lomonosov Ridge.
 76 Along a parallel drift track ca. 150 km closer to
 77 Greenland (Fig. 1), ice station Arlis-II traversed
 78 and looped over seafloor morphology interpreted



1 Fig. 2. Seismic reflection lines crossing the NP-28 Channel along the foot of the Lomonosov Ridge. Profile locations in Fig. 1
 2 (bold white lines). Arrows on seismic section NP-28 indicate past locations of the channel axis.

79 as a ca. 13-km-wide and more than 200-m-deep
 80 channel (Fig. 2). All channel crossings show a
 81 higher levee on the east side. The depth contours
 82 between 84°30'N and 89°N in the improved In-
 83 ternational Bathymetric Chart of the Arctic

Ocean (Jakobsson et al., 2001) clearly express
 the path of the channel and its associated levee
 towards the margin of North Greenland–Canada.
 The main channel is likely to be fed by tributary
 channels on the continental slope of North Green-
 84
 85
 86
 87
 88



1 Fig. 3. Line drawing of a multichannel seismic section across Amundsen Basin. Sequence nomenclature from Jokat et al. (1995).

89 land inferred from bathymetric depressions (Fig.
90 1).

91 In the Amundsen Basin, sequences AB-7 and
92 AB-8 (Jokat et al., 1995) exhibit upward convex
93 internal horizons with maximum thickness near
94 the North Pole (Fig. 3). Two submarine channels
95 are present at the seafloor culmination. The posi-
96 tion of the channels at the highest elevation of the
97 abyssal plain and internal units of the sequence,
98 particularly the lowermost one tapering off to-
99 wards either side, suggests that sequences AB-7
100 and AB-8 represent the distal lobe of a submarine
101 fan. The two channels at the seafloor are very
102 different in size and cross section; the largest
103 ones have a channel floor width of maximally
104 1.8 km, and a cross section with one levee higher
105 (35 m) than the other. A width-to-depth ratio less
106 than 40 is compatible with a location on the lower
107 part of a fan (Clark and Pickering, 1996). The
108 smaller channel is less than 0.4 km wide and the
109 difference in levee height is 5 m (Fig. 4). Small
110 bow-tie reflections within sequence AB-8 are in-
111 terpreted as relict channels (for example channel
112 X in Figs. 3 and 4). The low resolution in the data
113 makes it difficult to assess the degree of vertical
114 and lateral channel stacking in each case, apart
115 from noting that the channels have occupied dif-
116 ferent positions during fan growth. Most likely
117 the channels are to be classified as aggradational
118 and a low gradient would also imply likely high
119 sinuosity of the channel path (Clark et al., 1992).
120 The low relief of the smaller modern channel near
121 the base of the Lomonosov Ridge may reflect its
122 relative youthfulness.


123 A 12.9-m-long sediment core (PS2190; Fuetter-
124 er, 1992) was recovered about 6.5 km north of the
125 smaller channel (Figs. 1 and 3). This location is
126 actually 3.3 km from the geographical North
127 Pole. The upper 2 m of the core consist of clay
128 with alternating layers of dark brown and olive
129 colour interrupted by two sandy layers between 50
130 cm and 90 cm depth. Below 2 m subbottom is a
131 stack of silty sand layers, fining upwards. Most of
132 the turbidite layers are 10 cm thick (maximally 20
133 cm), and separated by mottled intervals of similar
134 thickness. The sediment stratigraphy shows that
135 the coring site has received deposition of hemipe-
136 lagic clays repeatedly interrupted by overbank

137 flow from turbidity current events in the smaller
138 channel. The sediments about 245 km farther into
139 the abyssal plain along longitude 110°E (core
140 PS2176; Fig. 1) consist of a 2-m-thick upper
141 clay interval overlying alternating intervals of
142 dark grey silt grading into clay, often laminated
143 at the base. Clay intervals in this 14-m-long core
144 are more dominant compared to the core
145 (PS2190) from the fan lobe. The site of core
146 PS2176 must be distal to any fan lobes extending
147 from either the Greenland or Siberian continental
148 margin (Fig. 1). The most important sediment
149 transport in this part of the basin probably occurs
150 in a near bottom nepheloid layer (Hunkins et al.,
151 1969) and as the vertical flux from fallout of bio-
152 genic matter and ice transported lithic material
153 (Clark and Hanson, 1983; Reimnitz et al., 1994).

3. Fan development and timing 154

155 Sequences AB-7 and AB-8 represent the emer-
156 gence of a submarine fan lobe extending from the
157 polar margin to the central part of the Amundsen
158 Basin and mark a change in mode and flux of
159 sediment input to the abyssal plain. The crest of
160 the distal depositional lobe appears to have main-
161 tained its position through time (Fig. 3) while at
162 least a section of the NP-28 Channel between
163 88°N and 89°N has migrated more than 50 km
164 towards the foot of the Lomonosov Ridge (Fig.
165 2). We suggest that the key elements in develop-
166 ment of the North Pole Submarine Fan are in-
167 creased sediment input to the margin of Canada
168 and North Greenland and topographic focussing
169 of gravity driven flows. The east–west trending
170 saddle between the southern end of the Lomonosov
171 Ridge and the Lincoln Sea margin collects
172 sediment input from a ca. 200-km-long section
173 of the margin into the NP-28 Channel (Fig. 1).
174 Apparent bathymetric indentations in the North
175 Greenland continental margin west and east of
176 Morris Jesup Rise may represent tributary path-
177 ways to the main trunk channel (Fig. 1). Turbid-
178 ity currents in the NP-28 Channel subject to the
179 Coriolis force would create the observed levee
180 configuration by over-bank flow (Fig. 2).

181 We have sparse stratigraphic information for



1 Fig. 4. Crossings of sediment pathways to the Amundsen Basin east of Morris Jesup Rise. Channel location numbers in Figs. 1
2 and 3.

182 the sediments in the Amundsen Basin. The Holo-
 183 cene sedimentation rate at the location of core PS
 184 2190 on the fan lobe (Fig. 1) is 1.4 cm/kyr or less
 185 (Gard, 1993). Schneider et al. (1996) attempted to
 186 date the same 12.9-m-long core by the palaeomag-
 187 netic method and concluded in spite of variable
 188 grouping of the inclinations that the sediments
 189 were probably deposited within the Brunhes nor-
 190 mal geomagnetic polarity interval (i.e. sedimenta-
 191 tion rate > 1.7 cm/kyr).

192 The maximum sediment input to the Amundsen
 193 Basin must be related to glacial maxima or degla-
 194 ciation events. Modeling suggest that the Arctic
 195 Ocean received about 15% of the total Northern
 196 Hemisphere iceberg flux during the Last Glacial
 197 Maximum, predominantly from North Canada
 198 (Bigg and Wadley, 2001). Field results demon-
 199 strate that the Nares Strait was completely filled
 200 with ice and transport from Greenland dominated
 201 the northern end of the strait where erratics were
 202 brought to more than 800 m above sea level, 5–10
 203 km inland on the Ellesmere coast (England,
 204 1999). Ice streams from this area must have en-
 205 tered the Lincoln Sea and brought sediments to
 206 the shelf edge where a depocentre is suggested by
 207 seaward convex bathymetric contours on the
 208 upper continental slope. Downslope gravity driv-
 209 en flows may have transported the material far-
 210 ther to the deep Amundsen Basin.

211 The earliest direct evidence of glaciation on the
 212 polar margin in the Southwest Canadian Arctic
 213 Archipelago on Banks Island postdates 1.77 Ma
 214 (Barendregt et al., 1998). Subsequently, at least
 215 two and possibly five full continental glaciations
 216 are recorded within the Matuyama reversed geo-
 217 magnetic polarity interval and three glaciations
 218 within the Bruhnes normal zone (< 0.78 Ma).
 219 Results of ODP drilling west of Svalbard date
 220 the first glacial advance to the shelf edge at ca.
 221 1.6 Ma, associated with onset of frequent diamic-
 222 tic debris flow events (Forsberg et al., 1999).
 223 However, input of ice rafted detritus north of
 224 Svalbard increased dramatically at about 2.5 Ma
 225 (Thiede et al., 1995) and large glacial fans formed
 226 along the margin west of Svalbard (Faleide et al.,
 227 1996) and also along the East Greenland conti-
 228 nental margin (Solheim et al., 1998). Intervals of
 229 debris flows at Site 987 off Scoresby Sound, East-

Central Greenland, suggest that glaciers first ex-
 230 tended to the shelf edge between 5 and 4.6 Ma,
 231 and later between 2.5–1.6 Ma (Channell et al.,
 232 1999; Solheim et al., 1998). Average sedimenta-
 233 tion rates at these ice proximal sites (Site 986,
 234 water depth 2050 m; Site 987, water depth 1680
 235 m) remained greater than 20 cm/kyr during the
 236 period 2.5–1 Ma (Channell et al., 1999). Although
 237 influx of glacial sediments to the Amundsen Basin
 238 from North Greenland and the Canadian Arctic
 239 may have started in the latest Miocene, we infer
 240 that deposition of most of sequences AB-7 and
 241 AB-8 is related to input from increased erosion
 242 during the full scale Northern Hemisphere glacia-
 243 tion after 2.5 Ma.
 244

4. Discussion and conclusions 245

246 The Amundsen Basin is a closed elongated ba-
 247 sin, and the principal terrestrial sediment input
 248 from the time of submergence of the Lomonosov
 249 Ridge must be from both ends of the basin; the
 250 margin north of the New Siberian Islands and the
 251 Greenland/Canadian Arctic continental margin.
 252 On the European polar margin, the evidence for
 253 a large Plio–Pleistocene sediment influx to the
 254 Nansen Basin from the Svalbard continental mar-
 255 gin is fairly well established (Rasmussen and
 256 Fjeldskaar, 1996; Vågnes, 1996; Thiede et al.,
 257 1995). A number of glacial through-mouth fans
 258 formed on the circum-arctic continental margin,
 259 but it is presently not known if any developed
 260 into submarine fans with a channel/levee system
 261 which extended far into the adjacent abyssal
 262 plain. Three seismic transects demonstrate that
 263 the gentle continental slope north of Greenland
 264 adjacent to the Lomonosov Ridge is constructed
 265 by levee deposits from the main channel of a sub-
 266 marine fan which extends beyond the North Pole
 267 in the Amundsen Basin. We suggest that the ex-
 268 istence of the North Pole Fan is partly due to the
 269 presence of the elevated southern end of Lomo-
 270 nosov Ridge which formed a sediment trap to
 271 downslope glacial debris flows for a distance of
 272 200 km along the Lincoln Sea margin. Large
 273 amounts of glacial debris accumulated in the
 274 deepest passage for further gravity driven trans-

275 port into the abyssal realm through the NP-28
276 Channel. During pre-glacial times, the continental
277 margins would have shed siliclastic debris into the
278 Amundsen Basin during sea-level lowstands and
279 the total sediment input would have been substan-
280 tially lower.

281 5. Uncited references

282 [Funder et al., 1985](#)

283 Acknowledgements

284 This paper is dedicated to the memory of Mi-
285 khail Y. Sorokin who was suddenly taken away
286 by a tragic event. We thank the crews of ice sta-
287 tions Arlis-II and NP-28, and the officers and
288 crew of icebreakers *Polarstern* and *Oden* for their
289 effort in collecting the data. The Russian ice sta-
290 tion NP-28 operated from 25 February 1987 until
291 17 January 1989 and drifted ca. 2800 km during
292 that period.


293 References

294 Bigg, G.R., Wadley, M.R., 2001. The origin and flux of ice-
295 bergs released into the Last Glacial Maximum Northern
296 Hemisphere oceans: The impact of ice-sheet topography.
297 *J. Quat. Sci.* 16, 565–573.
298 Burkhanov, V.F., 1957. Soviet Arctic research: Priroda 5, 21–
299 30 (translated by E.R. Hope). Directorate of Scientific In-
300 formation, DRB Canada.
301 Barendregt, R.W., Vincent, J.-S., Irving, E., Baker, J., 1998.
302 Magnetostratigraphy of Quaternary and late Tertiary sedi-
303 ments on Banks Island, Canadian Arctic Archipelago. *Can.*
304 *J. Earth Sci.* 35, 147–161.
305 Channell, J.E.T., Smelror, M., Jansen, E., Higgins, S.M., Leh-
306 man, B., Eidvin, T., Solheim, A., 1999. Age models for
307 glacial fan deposits off East Greenland and Svalbard (Sites
308 986 and 987). In: Jansen, E., Raymo, M.E., Blum, P., Her-
309 bert, T. (Eds.), Proceedings ODP, Scientific Results 162.
310 Ocean Drilling Program, College Station, TX.
311 Clark, J.D., Pickering, K.T., 1996. Submarine Channels: Pro-
312 cesses and Architecture. Vallis Press, London, 230 pp.
313 Clark, J.D., Kenyon, N.H., Pickering, K.T., 1992. Quantita-
314 tive analysis of the geometry of submarine channels: Impli-
315 cations for the classification of submarine fans. *Geology* 20,
316 633–636.
317 Clark, D., Hanson, A., 1983. Central Arctic Ocean sediment

texture: A key to ice transport mechanisms. In: Molnia, B. 318
(Ed.), *Glacial-marine Sedimentation*. Plenum, New York, 319
pp. 301–330. 320
Dietz, R.S., Shumway, G., 1961. Arctic Basin geomorphology. 321
Geol. Soc. Am. Bull. 72, 1319–1330. 322
England, J., 1999. Coalescent Greenland and Innuitian ice dur- 323
ing the Last Glacial Maximum: Revising the Quaternary of 324
the Canadian High Arctic. *Quat. Sci. Rev.* 18, 421–456. 325
Faleide, J.I., Solheim, A., Fiedler, A., Hjelstuen, B.O., Ander- 326
sen, E., Vanneste, K., 1996. Late Cenozoic evolution of the 327
western Barents Sea–Svalbard continental margin. *Glob.* 328
Planet. Change 12, 53–74. 329
Forsberg, C.F., Solheim, A., Elverhøi, A., Jansen, E., Chan- 330
nell, J.E.T., Andersen, E., 1999. The depositional environ- 331
ment of the western Svalbard margin during late Pliocene 332
and the Pleistocene: Sedimentary facies changes at Site 986. 333
In: Jansen, E., Raymo, M.E., Blum, P., Herbert, T. (Eds.), 334
Proceedings ODP, Scientific Results 162. Ocean Drilling 335
Program, College Station, TX. 336
Fuetterer, D.K., 1992. Arctic '91: The Expedition ARK-VIII/3 337
of RV 'Polarstern' in 1991. *Ber. Polarforsch.* 107, 267 pp. 338
Funder, S., Abrahamsen, N., Bennike, O., Freyling-Hansen, 339
R.W., 1985. Forested Arctic: Evidence from North Green- 340
land. *Geology* 13, 542–546. 341
Gard, G., 1993. Late Quaternary coccoliths at the North Pole: 342
Evidence of ice-free conditions and rapid sedimentation in 343
the central Arctic Ocean. *Geology* 21, 227–230. 344
Hunkins, K., Thorndike, E.M., Matieu, G., 1969. Nepheloid 345
layers and bottom currents in the Arctic Ocean. *J. Geophys.* 346
Res. 74, 6995–7008. 347
Jakobsson, M., and the IBCAO Editorial Board Members, 348
2001. Improvement to the International Bathymetric Map 349
of the Arctic Ocean (IBCAO): Updating the data base 350
and the grid model. Abstract Eos, *Trans. Am. Geophys.* 351
Union 84. 352
Jokat, W., Weigelt, E., Kristoffersen, Y., Rasmussen, T., 353
Schöne, 1995. New insights into the evolution of the Lomo- 354
nosov Ridge and the Eurasia Basin. *Geophys. J. Int.* 122, 355
378–392. 356
Ostenso, N., Wold, R.J., 1977. A seismic and gravity profile 357
across the Arctic Ocean. *Tectonophysics* 37, 1–24. 358
Rasmussen, E., Fjeldskaar, W., 1996. Quantification of the 359
Pliocene–Pleistocene erosion of the Barents Sea from present 360
day bathymetry. *Glob. Planet. Change* 12, 119–133. 361
Reimnitz, E., Dethleff, D., Nurnberg, D., 1994. Contrasts in 362
Arctic sea-ice regimes and some implications: Beaufort Sea 363
versus Laptev Sea. *Mar. Geol.* 119, 215–225. 364
Schneider, D.A., Backman, J., Possnert, G., 1996. Paleomag- 365
netic constraints on sedimentation rates in the Eastern Ar- 366
ctic Ocean. *Quat. Res.* 46, 62–71. 367
Solheim, A., Faleide, J.-I., Andersen, E., Elverhøi, A., Fors- 368
berg, C.F., Vanneste, K., Uenzelmann-Neben, G., Channell, 369
J.E.T., 1998. Late Cenozoic seismic stratigraphy and glacial 370
geological development of the East Greenland and Sval- 371
bard–Barents Sea continental margins. In: Elverhøi, A., 372
Dowdeswell, J., Funder, S., Mangerud, J., Stein, R. (Eds.), 373

- 374 Glacial and Oceanic History of the Polar North Atlantic 382
375 Margins. *Quat. Sci. Rev.* 17, 155–184. 383
376 Thiede, J., Myhre, A., Firth, J.V., and the Shipboard Scientific 384
377 party, 1995. Cenozoic Northern Hemisphere polar and sub- 385
378 polar ocean paleoenvironments (Summary of ODP Leg 151 386
379 drilling results). In: Myhre, A.M., Thiede, J., Firth, J.V., et 387
380 al., *Proc. Ocean Drilling Program, Initial Reports* 151, 397– 388
381 420.
- Vågnes, E., 1996. Cenozoic deposition in the Nansen Basin, a first order estimate based on present day bathymetry. *Glob. Planet. Change* 12, 149–157.
- Weber, J.R., 1983. Maps of the Arctic Basin sea floor: A history of bathymetry and its interpretation. *Arctic* 36, 121–142.

UNCORRECTED PROOF



-- trace of channel (NP-28 Channel) from IBCAO bathymetry